

EULER'S PARTITION THEOREM AND REFINEMENTS WITHOUT APPEAL TO INFINITE PRODUCTS

Krishnaswami Alladi

Dedicated to Peter Paule for his 60th Birthday

Abstract: *We present a new proof of Euler's celebrated theorem that the number of partitions of an integer into odd parts equals the number of partitions of that integer into distinct parts without using infinite product representations but only certain "amalgamation" properties and dissections of their series generating functions. Our approach also yields a simpler and more direct proof of a well known refinement of Euler's theorem due to Fine. We then use the same ideas along with conjugation of partitions into distinct parts to improve Fine's theorem and also to obtain a dual of a refinement of Euler's theorem due to Bessenrodt. Finally we discuss links between our approach and the Rogers-Fine identity.*

1. Introduction

One of the first results an entrant to the theory of partitions encounters is Euler's fundamental and beautiful theorem:

Theorem E: *Let $p_d(n)$ and $p_o(n)$ denote the number of partitions of n into distinct parts and odd parts respectively. Then*

$$p_d(n) = p_o(n).$$

Euler's proof of Theorem E made use of product representations of the generating functions of $p_d(n)$ and $p_o(n)$:

$$(1.1) \quad \sum_{n=0}^{\infty} p_d(n)q^n = \prod_{m=1}^{\infty} (1 + q^m) = \prod_{m=1}^{\infty} \frac{(1 - q^{2m})}{(1 - q^m)} = \prod_{m=1}^{\infty} \frac{1}{(1 - q^{2m-1})} = \sum_{n=0}^{\infty} p_o(n)q^n.$$

Euler's theorem and the simple yet fundamental idea in his proof, namely the replacement of expressions of the form $1+y$ by $(1-y^2)/(1-y)$ in products and the study of the resulting the cancellations, plays a crucial role in the theory of partitions (see Andrews [3], for instance). Many proofs of Euler's theorem are known and a variety of important refinements of it have been obtained by Sylvester [11], Fine [8], Bessenrodt [5] and others. Our approach here (see section 2) is to prove Euler's theorem by *only* considering the series generating function of $p_d(n)$ and an important (but under-utilized) *amalgamation property* of this series. We then convert the series generating function of $p_d(n)$ to the series generating function of $p_o(n)$ by a suitable *dissection* of the terms of the series. In doing so we use 2-modular Ferrers graphs to establish the equivalence. We then combine these ideas with the conjugation of the Ferrers graphs of partitions into distinct parts to improve a refinement of Euler's theorem due to Fine [8] and to obtain a dual of a refinement due to Bessenrodt [5].

2000 AMS Subject Classification: 05A17, 05A15, 05A19

Keywords: partitions, odd parts, distinct parts, Euler's theorem, series generating functions, amalgamation, even-odd split, 2-modular Ferrers graphs, refinement.

Sylvester [11] improved upon many partition theorems of Euler by combinatorial methods. In particular, using a graphical representation he was led to the Theorem S below which is a refinement of Theorem E. It is not easy to establish that the graphical representation yields a bijective proof of Theorem S; this was recently done by Kim and Yee [9].

Theorem S: *The number of partitions of an integer n into odd parts of which exactly k are different is equal to the number of partitions of n into distinct parts which can be grouped into k (maximal) blocks of consecutive integers.*

Yet another refinement of Theorem E was found by Fine, namely,

Theorem F: *Let $p_d(n; k)$ denote the number of partitions of n into distinct parts with largest part k . Let $p_o(n; k)$ denote the number of partitions of n into odd parts such that the largest part plus twice the number of parts is $2k+1$. Then*

$$p_d(n; k) = p_o(n; k).$$

Fine observed this in 1954 but published it only in his 1988 monograph [8]. Fine's proof of Theorem F was not combinatorial, but q -theoretic; it is sketched in [3]. Andrews ([3], p. 27) provides a q -theoretic proof in detail, but prior to that in 1966 [2], noted that Theorem F also falls out from Sylvester's graphical proof of Theorem E. Theorems S and F are *refinements* of Theorem E because by summing over k we get Theorem E. Our approach to Theorem E yields a much simpler and very direct proof of Theorem F and this is given in Section 3.

Bessenrodt [5] obtained the following elegant reformulation of Theorem E from Sylvester's bijection for Theorem S. This refinement is also a limiting case of the deep lecture hall partition refinement of Theorem E due to Bousquet-Melou and Eriksson [6].

Theorem B: *Let $p_{d,k}(n)$ denote the number of partitions of n into distinct parts such that the alternating sum starting with the largest part is k . Let $p_{o,k}(n)$ denote the number of partitions of n into odd parts with total number of parts equal to k . Then*

$$p_{d,k}(n) = p_{o,k}(n).$$

We will provide in Section 4 a simple direct proof of Theorem B by considering conjugates of the Ferrers graphs of partitions into distinct parts and then using the amalgamation-dissection ideas to convert the generating function of $p_{d,k}(n)$ into that of $p_{o,k}(n)$. This also yields a dual of Theorem B and an improvement of Theorem F (see Section 5).

As examples of some recent works pertaining to Theorem E (including its analogs and Glaisher's generalization to all odd moduli) and Theorem B, emphasizing combinatorial arguments, we mention the papers of Berkovich-Uncu [4], Straub [10], and of Xiong and Keith [12]. But our approach is quite different.

We shall use the standard notation

$$(1.2) \quad (a)_n = (a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j),$$

and

$$(1.3) \quad (a)_\infty = (a; q)_\infty = \lim_{n \rightarrow \infty} (a)_n = \prod_{j=0}^{\infty} (1 - aq^j), \quad \text{when } |q| < 1.$$

When the base is q , we write $(a)_n$ as in (1.2) for simplicity, but when the base is anything other than q , it will be displayed.

2. New proof of Theorem E

The series generating function of $p_d(n)$ is

$$(2.1) \quad \sum_{n=0}^{\infty} p_d(n)q^n = \sum_{m=0}^{\infty} \frac{q^{m(m+1)/2}}{(q)_m}.$$

This is due to Euler who noted that the generating function of all partitions having exactly m parts is $q^m/(q)_m$. From such partitions we get all partitions into m distinct parts by adding 0 to the smallest part, 1 to the second smallest part, ..., and $m - 1$ to the largest part. This procedure is reversible. So the term

$$(2.2) \quad \frac{q^{m(m+1)/2}}{(q)_m}$$

on the right in (2.1) is the generating function of partitions into distinct parts with exactly m parts.

The terms in (2.2) have an interesting *amalgamation property*, namely,

$$(2.3) \quad \frac{q^{(2m-1)2m/2}}{(q)_{2m-1}} + \frac{q^{2m(2m+1)/2}}{(q)_{2m}} = \frac{q^{2m^2-m}}{(q)_{2m}},$$

and its companion

$$(2.4) \quad \frac{q^{2m(2m+1)/2}}{(q)_{2m}} + \frac{q^{(2m+1)(2m+2)/2}}{(q)_{2m+1}} = \frac{q^{2m^2+m}}{(q)_{2m+1}}.$$

These amalgamation properties have not been fully exploited, and we shall use them here. In particular, we note from (2.1) and (2.3) that

$$(2.5) \quad \sum_{n=0}^{\infty} p_d(n)q^n = 1 + \sum_{m=1}^{\infty} \frac{q^{2m^2-m}}{(q)_{2m}}.$$

Next, having amalgamated consecutive pairs of terms in (2.1) to get (2.5), we will dissect the denominator terms in (2.5) into its odd and even components. More precisely, we rewrite (2.5) as

$$(2.6) \quad 1 + \sum_{m=1}^{\infty} \frac{q^{2m^2-m}}{(q)_{2m}} = 1 + \sum_{m=1}^{\infty} \frac{q^{2m^2-m}}{(q^2; q^2)_m (q; q^2)_m}.$$

We will now show that the series on the right of (2.6) is the generating function of $p_o(n)$.

Represent a partition into odd parts as a 2-modular Ferrers graph, namely a Ferrers graph in which there is a 1 at the node on the extreme right of each row, and there is a 2 at every other node. Consider the Durfee square in this 2-modular Ferrers graph, namely the largest square of nodes starting from the upper left hand corner. Let the Durfee square be

of dimension $m \times m$. Now the part below this Durfee square is a partition into odd parts the largest of which is $\leq 2m - 1$. The generating function of such partitions is

$$(2.7) \quad \frac{1}{(q; q^2)_m}.$$

The portion consisting of the Durfee square and the nodes to its right forms a partition into exactly m odd parts each $\geq 2m - 1$. If $2m - 1$ is removed from each of the m rows of this part of the graph, we remove $2m^2 - m$ in total. The remaining portion is a 2-modular Ferrer's graph with only twos in it, and which, if read columnwise, is a partition into even parts each $\leq 2m$. Thus the generating function of the Durfee square and portion to its right is

$$(2.8) \quad \frac{q^{2m^2-m}}{(q^2; q^2)_m}.$$

Thus we have shown that

$$(2.9) \quad 1 + \sum_{m=1}^{\infty} \frac{q^{2m^2-m}}{(q^2; q^2)_m (q; q^2)_m} = \sum_{n=0}^{\infty} p_o(n) q^n.$$

Theorem E follows from (2.9), (2.6) and (2.5).

3. Simple proof of Theorem F

The amalgamation-dissection idea of Section 2 yields the refined Theorem F as we show here. But first, we introduce the following notation for convenience: For any partition π , we let

$$\lambda(\pi) = \text{largest part of } \pi,$$

$$\nu(\pi) = \text{number of parts of } \pi,$$

and

$$\sigma(\pi) = \text{the sum of the parts of } \pi.$$

Thus $\sigma(\pi)$ is the integer being partitioned. Finally let D denote the set of partitions into distinct parts, and Ω , the set of partitions into odd parts.

By following the ideas of Euler that we described prior to (2.2), we get

$$(3.1) \quad \sum_{n=0}^{\infty} \frac{z^n t^n q^{n(n+1)/2}}{(tq)_n} = \sum_{\pi \in D} z^{\nu(\pi)} t^{\lambda(\pi)} q^{\sigma(\pi)}.$$

Even though this two parameter refined generating function of partitions into distinct parts is fundamental, it has not been given much attention because it does not have a product representation. However, if we set $t = 1$ and count only the number of parts, then we get a product representation for the expression in (3.1), namely

$$\prod_{k=0}^{\infty} (1 + zq^k),$$

and this product has been investigated in detail since the time of Euler. When we set $z = 1$ in (3.1) and keep track only of the largest part, we *do not* get a product representation for the series

$$(3.2) \quad \sum_{n=0}^{\infty} \frac{t^n q^{n(n+1)/2}}{(tq)_n} = \sum_{\pi \in D} t^{\lambda(\pi)} q^{\sigma(\pi)}.$$

Our emphasis here is on series and not infinite products. The terms of the series on the left in (3.2) amalgamate as in (2.3) and (2.4), and this is interesting and useful even though we do not have a product representation. More precisely we have,

$$(3.3) \quad \frac{t^{2m-1} q^{(2m-1)2m/2}}{(tq)_{2m-1}} + \frac{t^{2m} q^{2m(2m+1)/2}}{(tq)_{2m}} = \frac{t^{2m-1} q^{2m^2-m}}{(tq)_{2m}}.$$

and its companion

$$(3.4) \quad \frac{t^{2m} q^{2m(2m+1)/2}}{(tq)_{2m}} + \frac{t^{2m+1} q^{(2m+1)(2m+2)/2}}{(tq)_{2m+1}} = \frac{t^{2m} q^{2m^2+m}}{(tq)_{2m+1}}.$$

Thus using (3.3) we get

$$(3.5) \quad \sum_{n=0}^{\infty} \frac{t^n q^{n(n+1)/2}}{(tq)_n} = 1 + \sum_{m=1}^{\infty} \frac{t^{2m-1} q^{2m^2-m}}{(tq)_{2m}}.$$

As in (2.6), we dissect the denominator on the right in (3.5) and rewrite it as

$$(3.6) \quad \sum_{n=0}^{\infty} \frac{t^n q^{n(n+1)/2}}{(tq)_n} = 1 + \sum_{m=1}^{\infty} \frac{t^{2m-1} q^{2m^2-m}}{(tq^2; q^2)_m (tq; q^2)_m}.$$

We will now combinatorially interpret the coefficient of t^k in (3.6).

We know already from (2.7) and (2.8) that the expression

$$(3.7) \quad \frac{t^{2m-1} q^{2m^2-m}}{(tq^2; q^2)_m (tq; q^2)_m}$$

without the parameter t , is the generating function of partitions π^* into odd parts whose 2-modular Ferrers graphs have an $m \times m$ Durfee square. How are the powers of t generated in the expression in (3.7) and what does the power of t represent in relation to the partition π^* ? To understand this, we write the exponent k in the power of t , say t^k , in (3.7) as

$$(3.8) \quad k = (2m - 1) + i + j,$$

where t^i is generated from the factor $(tq^2; q^2)_m$, t^j is generated by the factor $(tq; q^2)_m$, and t^{2m-1} comes from the numerator. Thus from the arguments underlying (2.8), we see that in the 2-modular graphs under consideration, $(2m - 1) + 2i = \lambda(\pi^*)$, the largest part. Also $m + j = \nu(\pi^*)$, the number of parts, because there are j parts below the Durfee square, and the Durfee square is of size $m \times m$. Note that (3.8) yields

$$(3.9) \quad \lambda(\pi^*) + 2\nu(\pi^*) = (2m - 1) + 2i + 2(m + j) = 2k + 1.$$

On the other hand, the coefficient of t^k in (3.2) is the generating function of the number of partitions into distinct parts with largest part k . So by comparing coefficients of t^k on both sides of (3.6), we get Theorem R from (3.2) and (3.8).

Remark: As mentioned in the introduction, Fine's proof of Theorem F (see [8], p. 29 and p. 46) was not combinatorial but involved transformations of q -series. Andrews [3; p. 27] provides details of a q -theoretic proof of Theorem R using Heine's transformation and its consequences. Our proof given above is simpler and more direct because all that is needed is to amalgamate pairs of consecutive terms of the series generating function for partitions into distinct parts, and then to dissect the denominator of the resulting expression into its odd and even components. This actually leads to a refinement of Theorem F described below.

A further refinement:

Note that in the amalgamation in (2,3) and (3.3), we are adding the generating functions of partitions π into distinct parts for which the number of parts $\nu(\pi) = 2m - 1$ or $2m$. After the amalgamation and the dissection, we interpreted the expression in (3.7) as the generating function of partitions into odd parts π^* whose 2-modular Ferrer's graphs have an $m \times m$ Durfee square. Thus we have the the following refinement of Theorem F:

Theorem F*: *Let $p_d(n; k, m)$ denote the number of partitions π of n into distinct parts with $\lambda(\pi) = k$ and $\nu(\pi) = 2m - 1$ or $2m$. Let $p_o(n; k, m)$ denote the number of partitions π^* of n into odd parts with $\lambda(\pi^*) + 2\nu(\pi^*) = 2k + 1$ and such that the 2-modular Ferrers graph of π^* has a Durfee square of dimension $m \times m$. Then*

$$p_d(n; k, m) = p_o(n; k, m).$$

4. Conjugation of partitions into distinct parts

If π is a partition into distinct parts, then its conjugate π^* , namely the partition obtained by representing π as a Ferrers graph and considering the conjugate of this graph, is a partition whose set of parts is the set of consecutive integers from 1 up to $\nu(\pi)$. Thus letting ν denote $\nu(\pi)$, we have

$$(4.1) \quad \sigma(\pi) = \sigma(\pi^*) = \sum_{i=1}^{\nu} i f_i,$$

where f_i represents the frequency with which i occurs in π^* . We may rewrite the sum on the right in (4.1) as

$$(4.2) \quad \sigma(\pi^*) = (f_1 + f_2 + \dots + f_{\nu}) + (f_2 + f_3 + \dots + f_{\nu}) + \dots + (f_{\nu}),$$

where the quantities within the parenthesis represent the parts of π in decreasing order.

If we denote by $s(\pi)$ as in [5], the alternating sum of the parts of π starting with the largest part, then

$$(4.3) \quad \begin{aligned} s(\pi) &= (f_1 + f_2 + \dots + f_{\nu}) - (f_2 + f_3 + \dots + f_{\nu}) + (f_3 + f_4 + \dots + f_{\nu}) - (f_4 + f_5 + \dots + f_{\nu}) + \dots \\ &= f_1 + f_3 + f_5 + \dots =: \nu_o(\pi^*), \end{aligned}$$

where

$$\nu_o(\pi^*) = \text{the number of odd parts of } \pi^*.$$

So let us reformulate Theorem B as follows:

Theorem C: Let C denote the set of partitions with the property that all integers up to the largest part occur as parts. Let $p_C^*(n; \ell)$ denote the number of partitions π^* of n , $\pi^* \in C$, with $\nu_o(\pi^*) = \ell$.

Let $p_o^*(n; \ell)$ denote the number of partitions of n into odd parts such that the number of parts is ℓ . Then

$$p_C^*(n; \ell) = p_o^*(n; \ell)$$

A partition with the property that all integers up to the largest part occur as parts is known as a partition without gaps. We call such partitions as *chain partitions*, and C is the set of such partitions. Having reformulated Theorem B in terms of chain partitions, we give a simple proof of Theorem C.

Proof of Theorem C using series: For $\pi^* \in C$, consider its largest part $\lambda(\pi^*)$. If the largest part is $2j - 1$, then the generating function of such chain partitions π^* is

$$(4.4) \quad \frac{q^{2j^2-j} z^j}{(zq; q^2)_j (q^2; q^2)_{j-1}},$$

where the power of z in (4.4) is $\nu_o(\pi^*)$. If the largest part of π^* is $2j$, then the generating function is

$$(4.5) \quad \frac{q^{2j^2+j} z^j}{(zq; q^2)_j (q^2; q^2)_j}.$$

Now if we add the expressions in (4.4) and (4.5), they amalgamate to

$$\frac{q^{2j^2-j} z^j}{(zq; q^2)_j (q^2; q^2)_j}.$$

So we get

$$(4.6) \quad \sum p_C(n; \ell) z^\ell t^n = \sum_{j=0}^{\infty} \frac{q^{2j^2-j} z^j}{(zq; q^2)_j (q^2; q^2)_j}.$$

Just as we showed (2.9) via Durfee squares, it follows that the series on the right in (4.6) is

$$(4.7) \quad \sum_{j=0}^{\infty} \frac{q^{2j^2-j} z^j}{(zq; q^2)_j (q^2; q^2)_j} = \sum_n \sum_\ell p_o^*(n; \ell) z^\ell q^n.$$

Theorem C follows from (4.6) and (4.7) without any appeal to infinite products.

Graphical proof of Theorem C: We now provide a bijective proof of Theorem C using 2-modular graphs.

Start by representing a partition π into odd parts as a 2-modular graph. We illustrate our bijective proof by considering the partition $25+21+15+15+13+9+7+7+7+3+1+1+$. In this 2-modular graph, mark out the Durfee square. This is illustrated in Fig. 1.

Next delete the right most column of the Durfee square, fill them with ones, and move any twos that were in the right most column to the extreme right position on the same row. Thus the integer entries in the modified Durfee square (say of dimension j) add up to $2j^2 - j$. Group the integer entries in the modified Durfee square as indicated in Fig 2 to see that these represent the integers $1, 2, \dots, 2j - 1$. The rows below the modified square represent odd parts $\leq 2j - 1$. The columns to the right of the modified square represent even parts $\leq 2j$. Thus if the modified graph is viewed in this fashion, we get a chain partition π^* with largest part either $2j - 1$ or $2j$. Notice that the number of parts of π equals then number of odd parts of π^* and this proves Theorem C.

Fig 1: 5 x 5 Durfee Square





Fig 2:



5. A dual of Theorem C and an improvement of Theorem F

The graphical proof of Theorem C has interesting implications.

In the graphical proof given above, we focused on the number of parts. We now see what happens if we consider the largest part.

Suppose the size of the largest part of the partition π represented in Figure 1 is $2k + 1$. As before we let the size of the Durfee square and the modified Durfee square to be j . This means that the largest odd part $\lambda_o(\pi^*)$ of the chain partition π^* given by Fig. 2 is $2j - 1$. Consequently, $2, 4, \dots, 2j - 2$ occur as even parts of π^* and these account for $j - 1$ even parts. The number of even parts of π^* is given by $j - 1$ plus the number of columns to the right of the modified Durfee square. We may interpret $j - 1$ as the number of twos in the first row of the modified Durfee square in Fig. 2. Thus the number of even parts of π^* is k . This leads to the following *dual* of Theorem C:

Theorem C*:

Let $p_{o,k}^*(n)$ denote the number of partitions of n into odd parts with largest part $2k + 1$.
Let $p_{C,k}^*(n)$ denote the number of chain partitions of n with k even parts. Then

$$p_{o,k}^*(n) = p_{C,k}^*(n)$$

Remarks:

(i) Theorem C* is a dual of Theorem C because in Theorem C we counted the number of odd parts of π whereas in Theorem C* we count the size of the largest odd part of π ; similarly in Theorem C we count the number of odd parts of π^* whereas in Theorem C* we count the number of even parts of π^* . Thus by reformulating Theorem B in terms of Theorem C, we have arrived at the dual Theorem C*.

(ii) By combining Theorems C and C* we get Fine's Theorem F. This is because with $\nu_e(\pi^*)$ representing the number of even parts of π^* , we can write Theorem C* in the form

$$(5.1) \quad \lambda(\pi) = 2\nu_e(\pi^*) + 1.$$

Similarly we may write Theorem C in the form

$$(5.2) \quad \nu(\pi) = \nu_o(\pi^*).$$

Thus (5.1) and (5.2) yield

$$(5.3) \quad \lambda(\pi) + 2\nu(\pi) = 2\nu_e(\pi^*) + 1 + \nu_o(\pi^*) = 2\nu(\pi^*) + 1.$$

By taking the conjugate of π^* the number of parts of π^* is converted to the largest part of the conjugate partition, which is a partition into distinct parts, and this is precisely Fine's theorem. Thus Theorems C and C* are improvements of Fine's theorem.

(iii) In section 4 we noted that the infinite series in (4.6) and (4.7) can be interpreted in two different ways to be realized as the generating function of the two partition functions $p_C^*(n; \ell)$ and $p_o^*(n; \ell)$ in Theorem C. Similarly, the analytic version of Theorem C* is

$$(5.4) \quad \sum_n \sum_k p_{o,k}^*(n) w^k q^n = 1 + \sum_{k=0}^{\infty} \frac{w^k q^{2k+1}}{(q; q^2)_{k+1}} = 1 + \sum_{j=1}^{\infty} \frac{w^{j-1} q^{2j^2-j}}{(q; q^2)_j (wq^2; q^2)_j} = \sum_n \sum_k p_{C,k}^*(n) w^k q^n$$

Note that the series on the right in (5.4) is the dual of the series in (4.6) and (4.7) because in (4.6) the power of z is counting the number of odd parts, whereas on the right in (5.4) the power of w is counting the number of even parts.

Identities (4.6), (4.7), and (5.4) can be combined into a single identity as follows:

$$(5.5) \quad 1 + \sum_{k=0}^{\infty} \frac{w^k z q^{2k+1}}{(zq; q^2)_{2k+1}} = 1 + \sum_{j=1}^{\infty} \frac{w^{j-1} z^j q^{2j^2-j}}{(zq; q^2)_j (wq^2; q^2)_j}.$$

Note that in (4.6) and (4.7) we did not have the series on the left in (5.5) with $w = 1$ because we did not need it. Instead we interpreted the series on the right in (5.5) with $w = 1$ in two different ways one of which relied on an amalgamation.

It turns out that the series on the right in (5.5) is a special case of a certain variant of the Rogers-Fine identity as we shall see in the next section.

6. Connection with the Rogers-Fine identity

In the previous section we studied the 2-modular graphs of partitions into odd parts by keeping track of the largest part and the number of parts. What if we also keep track of the number of different odd parts? This leads us to a variant of the Rogers-Fine identity as we show now.

The Rogers-Fine identity in the form obtained by Fine [8; 14.1] is

$$(6.1) \quad F(\alpha, \beta, \tau; q) =: \sum_{n=0}^{\infty} \frac{(\alpha q)_n \tau^n}{(\beta q)_n} = \sum_{n=0}^{\infty} \frac{(\alpha q)_n (\alpha \tau q / \beta)_n \beta^n \tau^n q^{n^2} (1 - \alpha \tau q^{2n+1})}{(\beta q)_n (\tau)_{n+1}}.$$

Fine proved it by considering transformation properties of $F(\alpha, \beta, \tau; q)$ defined by the series on the left in (6.1).

In [1] we obtained the following variant of the Rogers-Fine identity:

$$(6.2) \quad f(a, b, c; q) =: 1 + \sum_{k=1}^{\infty} \frac{(1-a)(abq)_{k-1} b c^k q^k}{(bq)_k} = 1 + \sum_{j=1}^{\infty} \frac{b^j c^j q^{j^2} (1-a)(abq)_{j-1} (acq)_{j-1} (1-abcq^{2j})}{(bq)_j (cq)_j}.$$

Fine's function f and our function f are connected by the relation

$$(6.3) \quad \frac{(1-bq)}{(1-a)bcq} \{f(a, b, c; q) - 1\} = F(ab, bq, cq; q)$$

and so (6.1) and (6.2) are equivalent. The reason we investigated (6.2) was because it is combinatorially more interesting, and also can be established combinatorially in a very direct fashion.

The function $f(a, b, c; q)$ defined by the series on the left in (6.2) is the generating function of unrestricted partitions in which the power of b keeps track of the number of parts, the power of c keeps track of the largest part, and the power of $1-a$ keeps track of the number of different parts. It is to be noted that for unrestricted partitions the generating function $f(a, b, c; q)$ has an infinite product representation only when b or c equals 1. Thus when one keeps track of all three statistics $\lambda(\pi)$, $\nu(\pi)$, and $\nu_d(\pi)$ (= the number of different parts

of π), then one will NOT have a product representation but will have to deal only with a series representation. This is in line with the philosophy of this paper emphasizing series and removing dependence on infinite product representations. In order to pass from the defining series of f to the series on the right in (6.2), we studied in [1] the Ferrers graphs of unrestricted partitions using Durfee squares and the fact that under conjugation $\lambda(\pi)$ and $\nu(\pi)$ get interchanged, and $\nu_d(\pi)$ remains invariant. We needed to use the invariance of $\nu_d(\pi)$ under conjugation only on the portion of the Ferrers graph to the right of the Durfee square. This aspect will be crucial in the remark below.

The ideas in [1] can be applied to the 2-modular Ferrers graphs of partitions into odd parts. Without getting into details, we simply point out that what this means is to replace

$$(6.4) \quad q \mapsto q^2, \quad \text{and} \quad b \mapsto bq^{-1}$$

in (6.2). This yields

$$(6.5) \quad \sum_{k=1}^{\infty} \frac{(1-a)(abq; q^2)_{k-1} b c^{k-1} q^{2k-1}}{(bq; q^2)_k} = \sum_{j=1}^{\infty} \frac{b^j c^{j-1} q^{2j^2-j} (1-a)(abq; q^2)_{j-1} (acq; q^2)_{j-1} (1-abcq^{4j-1})}{(bq; q^2)_j (cq^2; q^2)_j}.$$

Note that if we set $a = 0$, that is if we do not count the number of different odd parts in the graph, then (6.5) reduces to (5.5) with the identifications $b = z$ and $c = w$.

We conclude by showing how Sylvester's theorem can be deduced from (6.5). For this purpose we state the dual of Sylvester's theorem by replacing partitions into distinct parts by chain partitions. Under conjugation, given a partition π into distinct parts having k maximal blocks of consecutive integers, its conjugate, namely the chain partition π^* will have $k - 1$ parts less than the largest part that repeat. Thus the dual of Sylvester's theorem is:

Theorem S*: *The number of partitions of an integer into odd parts of which exactly k are different equals the number of chain partitions of that integer having $k-1$ parts less than the largest part that repeat.*

Theorem S* can be deduced from (6.5) as follows. First decompose

$$(6.6) \quad 1 - abcq^{4j-1} = cq^{2j}(1 - abq^{2j-1}) + (1 - cq^{2j}).$$

When $1 - abcq^{4j-1}$ is first replaced by $cq^{2j}(1 - abq^{2j-1})$ on the right in (6.5), the resulting series can be interpreted as the generating function of chain partitions with largest part even. Similarly when $1 - abcq^{4j-1}$ is next replaced by $(1 - cq^{2j})$ on the right in (6.5), the resulting series can be interpreted as the generating function of chain partitions with largest part odd. Thus Theorem S* will fall out of (6.5) and (6.6).

Remarks:

(i) In the case of Ferrers graphs of unrestricted partitions, the number of different parts is the number of corners and this is invariant under conjugation. When one considers the 2-modular graphs of partitions into odd parts, the number of different parts is the number of corners, but under conjugation we do not have a 2-modular graph. This awkwardness is circumvented by replacing the graph in Figure 1 by the graph in Figure 2 so that the portion to the right of the Durfee square consists only of twos. The number of different odd parts that are at least as large as $2j - 1$ where j is the dimension of the Durfee square is given

by the number of corners to the right of the Durfee square and we can keep track of this by conjugation since that portion of the graph in Figure 2 has only twos in it.

(ii) Zeng [10] has studied combinatorially the original Rogers-Fine identity (6.1) under the dilation $q \mapsto q^2$, and with α, β chosen suitably so as to deal with partitions into odd parts. Our approach uses the variant (6.2) and so is combinatorially more direct. Also we have preferred to replace partitions into distinct parts by chain partitions. Hence there are essential differences between our approach and Zeng's.

Acknowledgement: I thank the referee for a very careful reading of the manuscript and for helpful suggestions. I also thank Frank Garvan for help with the figures.

References

- 1) K. Alladi, "A new combinatorial study of the Rogers-Fine identity and a related theta series", *Int'l J. Num. Th.*, **5** (2009), 1311-1320.
- 2) G. E. Andrews, "On basic hypergeometric series, mock theta functions, and partitions - II", *Quart. J. Math., Oxford*, **17** (1966), 132-143.
- 3) G. E. Andrews, "The theory of partitions", *Encyclopedia of Math. and its Appl.*, **2**, Addison-Wesley, Reading (1976).
- 4) A. Berkovich and A. K. Uncu, "On partitions with a fixed number of even-indexed and odd-indexed odd parts", *J. Num. Th.*, **167** (2016), 7-30.
- 5) C. Bessenrodt, "A bijection for Lebesgue's partition identity in the spirit of Sylvester", *Discrete Math.* **132** (1994), 1-10.
- 6) M. Bousquet-Melou and K. Eriksson, "Lecture hall partitions", *Ramanujan J.*, **1** (1997), 101-111.
- 7) W. Y. C. Chen, H. Y. Gao, K. Q. Ji, and M. Y. X. Li, "A unification of two refinements of Euler's partition theorem", *Ramanujan J.*, **23** (2010), 137-149.
- 8) N. J. Fine, "Basic hypergeometric series and applications", *Math. Surveys and Monographs*, **27** Amer. Math. Soc., Providence (1988).
- 9) D. Kim and A. J. Yee, "A note on partitions into distinct parts and odd parts", *Ramanujan J.*, **3** (1999), 227-231.
- 10) A. Straub, "Core partitions into distinct parts and an analog of Euler's theorem", *Europ. J. Comb.* **57** (2016), 40-49.
- 11) J. J. Sylvester, "A constructive theory of partitions arranged in three Acts, an Interact and an Exodion", *Amer. J. Math.*, **5** (1882), 251-330.
- 12) X. Xiong and W. Keith, "Euler's partition theorem for all moduli and new companions to the Rogers-Ramanujan-Andrews-Gordon identities", *Ramanujan J.* (to appear).
- 13) J. Zeng, "The q -variations of Sylvester's bijection between odd and strict partitions", *Ramanujan J.* **9** (2005), 289-303.

Department of Mathematics
University of Florida
Gainesville, FL 32611, USA
e-mail: alladik(at)ufl.edu